

Israel

2010 total population: 7 418 400

Income group: High

NCD mortality

2008 estimates	males	females
Total NCD deaths (000s)	15.6	16.6
NCD deaths under age 60 (percent of all NCD deaths)	17.4	10.7
<i>Age-standardized death rate per 100 000</i>		
All NCDs	376.5	267.9
Cancers	131.5	101.4
Chronic respiratory diseases	24.8	15.2
Cardiovascular diseases and diabetes	138.9	93.8

Behavioural risk factors


2008 estimated prevalence (%)	males	females	total
Current daily tobacco smoking	23.9	14.9	19.3
Physical inactivity

Metabolic risk factors





2008 estimated prevalence (%)	males	females	total
Raised blood pressure	38.3	33.5	35.8
Raised blood glucose	10.4	9.6	10.0
Overweight	62.4	59.4	60.9
Obesity	23.2	29.0	26.2
Raised cholesterol	51.5	55.5	53.6

Metabolic risk factor trends

Proportional mortality (% of total deaths, all ages)



NCDs are estimated to account for 87% of all deaths.



Country capacity to address and respond to NCDs

Has a Unit / Branch / Dept in MOH with responsibility for NCDs	No	Has an integrated or topic-specific policy / programme / action plan which is currently operational for:	
<i>There is funding available for:</i>		Cardiovascular diseases	No
NCD treatment and control	Yes	Cancer	No
NCD prevention and health promotion	Yes	Chronic respiratory diseases	DK
NCD surveillance, monitoring and evaluation	Yes	Diabetes	No
<i>National health reporting system includes:</i>		Alcohol	No
NCD cause-specific mortality	Yes	Unhealthy diet / Overweight / Obesity	No
NCD morbidity	Yes	Physical inactivity	No
NCD risk factors	DK	Tobacco	No
Has a national, population-based cancer registry	Yes	Number of tobacco (m)POWER measures implemented at the highest level of achievement	2/5

... = no data available

DK = Country responded "don't know"